

Summary of Wound Prevalence



Chronic Wounds




Population of Europe as of 28th Jan 2025 based on UN figures is:

- 744,687, 041 = 9.21% of world population.
- Chronic wounds have a prevalence of 2.21 per 1000.
- 24% of the population in a community area were in receipt of care by community nurses.

Venous Leg Ulcers


- **Approximately 20% of the population of Europe [28 countries]. are over 65 years = 101.1 million**
- **A point prevalence of .33 - 1.01 % has been reported.**



Percentage of Population over 65 Years (%)

- >23%
- 21.1-23
- 19.1-21
- 17.1-19
- 15-17
- <15%

Source: WHO, country data on diabetes.




Diabetic Foot Ulcers

Age-standardized prevalence of diabetes, 2016 (%)

- 10,0 or more
- 9,1 to 9,9
- 8,1 to 9,0
- 7,1 to 8,0
- 6,1 to 7,0
- 6,0 or less

Source: WHO, country data on diabetes.




- **Diabetes is one of the most common chronic conditions in Europe. At least 64 million adults and around 300 000 children and adolescents are estimated to be living with diabetes in the WHO European Region.**
- Regarding diabetic foot disease, it is estimated that between 19–34% of people with diabetes will develop this throughout their lifetime.
- Thus, if 64,300,000 have diabetes and approximately 25% will get a foot ulcer over their lifetime this is = to 16,075,000 in Europe.

Pressure ulcers/injuries

3.2 % - 32.4% of people in long-term care settings have a pressure ulcer [based on 20 studies of which 15 were from Europe].

Median prevalence of pressure ulcers in Europe was 10.8% [based on 79 articles] across all care settings.




Costs of Wound Care

Europe is the second largest region in the global wound care market.

The global market is projected to grow from USD 18.51 billion in 2022 to USD 30.04 billion by 2030.

Source: <https://www.fortunebusinessinsights.com/europe-wound-care-market-107613>

Prevalence of Chronic Wounds in Selective European Countries, 2022



References

1. Martinengo, L., et al., Prevalence of chronic wounds in the general population: systematic review and meta-analysis of observational studies. *Annals of Epidemiology*, 2019. 29: p. 8-15.
2. Skerritt, L., M. Gooney, and L. Sheahan, The provision of wound care and management in a community healthcare setting: an exploratory study. *J Wound Care*, 2025. 34(1): p. 74-87.
3. Zhang, P., et al., Global epidemiology of diabetic foot ulceration: a systematic review and meta-analysis (dagger). *Ann Med*, 2017. 49(2): p. 106-116.
4. Zhang, Y., et al., Global Disability Burdens of Diabetes-Related Lower-Extremity Complications in 1990 and 2016. *Diabetes Care*, 2020. 43(5): p. 964-974.
5. Probst, S., et al., Prevalence and incidence of venous leg ulcers-A systematic review and meta-analysis. *Int Wound J*, 2023. 20(9): p. 3906-3921.
6. Anthony, D., D. Alosoumi, and R. Safari, Prevalence of pressure ulcers in long-term care: a global review. *J Wound Care*, 2019. 28(11): p. 702-709.
7. Moore, Z., et al., The prevalence of pressure ulcers in Europe, what does the European data tell us: a systematic review. *J Wound Care*, 2019. 28(11): p. 710-719.
8. Zhang, H., et al., Incidence and prevalence of pressure injuries in children patients: A systematic review and meta-analysis. *J Tissue Viability*, 2022. 31(1): p. 142-151.
9. Jiao, Y., et al., Incidence of pressure injuries in fracture patients: A systematic review and meta-analysis. *J Tissue Viability*, 2022. 31(4): p. 726-734.